

Explosions inside Ejecta and Most Luminous Supernovae

S.I.Blinnikov

*Institute for Theoretical and Experimental Physics (ITEP), Moscow, 117218, Russia
RESCEU, School of Science,
Tokyo University, Tokyo 113-0033, Japan*

Abstract. The extremely luminous supernova SN2006gy is explained in the same way as other SNIIn events: light is produced by a radiative shock propagating in a dense circumstellar envelope formed by a previous weak explosion. The problems in the theory and observations of multiple-explosion SNe IIn are briefly reviewed.

Keywords: supernovae, explosion mechanism

PACS: 26.30.-k, 26.30.Ef, 97.60.Bw

INTRODUCTION

The discovery of SN2006gy [1, 2] demonstrates that some supernova (SN) events produce 10 or even 100 times more visible photons than other, already powerful explosions. The anomalously high power of the emission of SN2006gy demands an explanation.

SN2006gy is of SNIIn type and it revived interest in SN models where light is produced by a long living radiative shock which propagates in a dense circumstellar envelope.


I discuss problems in the theory of SNIIn events and in the physics of supercritical radiative shocks. The powerful visible light of SN2006gy can be easily explained by a radiative shock born due to a collision of SN ejecta with a dense cloud formed by a weak explosion some years before the SN. Strong X-ray emission of SNIIn near maximum light may be absent since it is absorbed by the dense cloud or not produced at all in the radiation-dominated shock.

SUPERNOVA TYPES


The models with a long living radiative shock running in a dense circumstellar envelope were invoked earlier [3, 4] to explain the unusual properties of other powerful supernovae with narrow emission lines in their spectra, SNIIn. SN2006gy also belongs to the same SNIIn class.

A simple diagram below illustrates the relation between different astronomical types of supernovae which are classified purely on the appearance of their spectra near maximum light, irrespective of the underlying physics. For example, we believe that SNe II are born when a giant star with an H-rich atmosphere has a powerful explosion in its core. This explosion may be a result of a catastrophic collapse of the stellar core. The


details of the explosion of core-collapsing SNe are still not clear, in spite of many important results obtained by workers in this field.



Taking into account the physics of SN explosion we can produce a couple of other, a bit different, diagrams for SN types. I want to subdivide the SN physics into two groups of problems. First, here is a diagram for the mechanism of explosion.



And the next diagram illustrates main ways to produce light during supernova explosions. Here S is entropy, and 'heating' is entropy production.



Remarkably, the details of the explosion are not important for explaining the light curves of many supernovae that are born from giant stars that retain their huge hydrogen envelopes. Successful SNII light models were already constructed four decades ago (largely by the work of the Soviet group in Moscow and Riga [5]) due to this insensitivity

TABLE 1. Four kinds of deaths for non-rotating stars

Main Seq. Mass	He Core	Supernova Mechanism
$10 \leq M \leq 95$	$2 \leq M \leq 40$	Fe core collapse to a neutron star or a black hole
$95 < M \leq 130$	$40 < M \leq 60$	Pulsational pair instability followed by Fe core collapse
$130 < M \leq 260$	$60 < M \leq 137$	Pair instability supernova
$260 < M$	$137 < M$	Black hole. Possible GRB ?

to details. The light of SNe II is the manifestation of entropy produced during a short period (hours to days) of the shock propagation in the body of the presupernova star. We cannot exclude the possibility that in some rare cases the SNe II are produced by thermonuclear explosions, not by a collapse, inside hydrogen envelopes. This may be the case for SNe IIn, and especially, SN2006gy. A supernova of type II shines for several months thanks to the heat stored in its body (the shock dies quickly), while in an SNIIn the heat (entropy) is replenished by the shock living several months.

If the hydrogen is lost by a massive star, then we have an SN Ib/c. The same (still unknown in details) core-collapse mechanism may lead to their explosions, as in SNe II, however, the light is due now to entropy produced by radioactivity: the decays $^{56}\text{Ni} \rightarrow ^{56}\text{Co} \rightarrow ^{56}\text{Fe}$ which lead to a slower heating of ejecta. This way of producing light is most important for SNe I of all subtypes. However, some contribution of radioactivity is clearly present in late light curves of type II supernovae as well, and for SN1987A in LMC it was important already before its maximum light, a month after the explosion.

If the radioactive mechanism was responsible for the light of SN2006gy, then the amount of ^{56}Ni must be higher than $10M_{\odot}$ [6, 7, 8]. This immediately implies a very large mass for the presupernova star, more than $100M_{\odot}$. More important, this implies a huge explosion energy, $(50 - 80) \times 10^{51}$ erg. One can delineate four kinds of deaths for massive stars, see Table 1. There is some uncertainty about the exact values due to uncertainties in mass-loss, rotation etc. Such massive stars do exist, and they can have powerful explosions in their oxygen cores which experience instability due to the creation of large numbers of electron-positron pairs.

PAIR INSTABILITY SUPERNOVAE

The word ‘instability’ refers here to hydrodynamics, to mechanical equilibrium, not to the process of pair creation which is quite stable and reversible in a thermodynamic sense in stellar interiors. A massive star loses its mechanical stability when pairs are being created because the adiabatic exponent γ goes down at $T \sim 0.1$ MeV, see Fig. 1: the work of contraction is spent in creating new particles and not for raising the momenta of particles that already exist and which provide for the equilibrium pressure.

We can easily estimate the path that leads the star into the domain of the pair-creation instability. From the virial theorem, for a star of mass M and radius R , omitting all coefficients of order unity, $P_c V \sim P_c R^3 \sim G_N M^2 / R$. Hence, the pressure P_c in the center is $P_c \simeq G_N M^2 / R^4$, while the density $\rho_c \simeq M / R^3$, and they are related as $P_c \simeq G_N M^{2/3} \rho_c^{4/3}$. So, if we have a classical ideal plasma with $P = \mathcal{R} \rho T / \mu$, where $\mathcal{R}$ is the universal gas




FIGURE 1. Left: adiabatic exponent in the low density asymptotic limit taking into account pair creation [9], see also [10]. Right: evolution track for one of the models [11] with initial $M \sim 103 M_{\odot}$ (solid line). The approximate boundary of pair-instability is shown by the dashed line

constant, and μ – mean molecular mass, we find

$$T_c \simeq G_N M^{2/3} \rho_c^{1/3} \mu / \mathcal{R}.$$

If we have a relativistic addition of aT^4 to P , the law $T_c \propto \rho_c^{1/3}$ is the same (but the coefficient is a bit different). These relations have been already shown at this conference by Naoki Yoshida and by Marco Limongi. The right panel of Fig. 1 shows how a massive star follows the law $T_c \propto \rho_c^{1/3}$ until entering the domain of pair-instability. We had already computed the light curves of pair-instability supernovae for $M > 130 M_{\odot}$ (third line in Table 1) some years ago (with S.Woosley and A.Heger), but we do not like them in the case of SN2006gy because we do not see the evidence for a tremendously high explosion energy in that case. If the explosion energy were 2 orders of magnitude higher than for normal SNe, then it should be seen in very broad spectral lines. What we see in SN2006gy is different: it has narrow P Cyg lines (hence, it is type IIn) superimposed on moderately broad emission component ($\sim 5 \times 10^3$ km/s [2]). There is no sign of huge kinetic energy in this event.

Multiple ejections in SNIIn

Supernovae of type IIn are among the most powerful transients in visible light. No radioactive material is needed to explain their light during the first several months: the light is produced by a long living radiative shock in a dense circumstellar envelope. This is the main difference with standard SNe II where the shock breaks out into rarefied interstellar medium and disappears quickly. Spectra and light curves of SNIIn can be explained only when the number density of circumstellar matter at radii of $\sim 10^{15-16}$ cm (where the narrow lines are formed) is unusually high, like $10^9-10^{10} \text{ cm}^{-3}$, see Fig. 2. This

implies that a large mass (on the order of $M_\odot$ and larger) must be ejected within years, or even months, before the observed SN. The first ejection may have kinetic energy appreciably lower than a standard supernova. The slow motion of its matter explains the narrow lines of a type IIn SN. The supernova itself is an explosion of a normal energy, but inside a dense cloud.

Paper [3] was the first to suggest that an SNIIn had a precursor, a relatively weak explosion ejecting a large slowly moving mass. A dramatically high mass loss needed for the formation of a dense envelope shortly before SNIIn 1994W (Fig. 2) has been derived in [4]. SNIIn 1995G is explained in [12] in a model similar to the one presented in [4]. Double explosions may be observed also for other SN types [13].

According to [14] an $\sim 11M_\odot$ star might produce strong flashes in the semi-degenerate O-Ne-Mg core a few years prior to SN explosion and the strongest flash could eject most of the hydrogen envelope with velocities ~ 100 km/s. More recent evolutionary computations do not support the conclusion on strong Ne flashes, but this complicated problem deserves further investigation.

Pulsational pair-instability

Another workable mechanism for multiple-explosion SNe has been proposed in our paper [11] to explain SN2006gy. It works for a high initial mass of the presupernova star $\sim 110M_\odot$ (see Fig. 2). This mechanism is also based on the pair-creation instability, but there is no catastrophic collapse or full explosion of the star.

The second line of Table 1 shows that between 95 and $130M_\odot$ a relatively unexplored phenomenon of *pulsational* pair instability supernova [15, 16, 17] occurs. An instability in the mechanical equilibrium is encountered, as in the heavier stars, during the evolution along $T_c \propto \rho_c^{1/3}$ path (Fig. 1). A thermonuclear explosion of oxygen occurs, but the energy released is inadequate to unbind the entire star. It suffices, however, to eject many solar masses of surface material, including the hydrogen envelope, in a series of giant ‘pulses’ (explosions of various strength).

The binding energy for the hydrogen envelope of $(95 - 130)M_\odot$ stars is $\sim (0.1 - 1) \times 10^{49}$ erg while the energy of an explosive pulse is about two orders of magnitude higher. The velocity is in the range $100 \div 5000$ km/s depending on amount of explosive burning and mass ejected. After a pulse, the remaining core contracts searching for a new equilibrium state. It obeys the $T_c \propto \rho_c^{1/3}$ law again, but now the mass is lower, so the track is a bit different (see Fig. 1). The time required for the contraction is sensitive to the strength of the first pulse. If it cools down severely after the pulse expansion, it may be centuries before the star ignites burning again. If it remains hotter than 1.5×10^9 K, it may only take days. After one, two or several explosion pulses the remnant of the massive star continues to live, contrary to other supernovae, and eventually it should collapse.

To explain SN2006gy, we consider the evolution of a star with initial mass $110M_\odot$. The evolution is calculated using the Kepler code [18, 6] with reduced mass loss. The star encounters pair-instability with a total mass of $75M_\odot$ (a helium core of $50M_\odot$) and experiences the first pulse ejecting a cloud with mass $\approx 25M_\odot$ and $E_{\text{kin}} \approx 1.4 \times 10^{50}$ erg.




FIGURE 2. Left: SN1994W structure [4] prototypical for SNe IIn. Right: light curve models for SN2006gy. Dots – observations [2], the last observed point is from [21], courtesy M.Tanaka. The solid line is the model discussed in the text (with numerical resolution higher than in [11]), and the dashed one where the velocity of all the ejecta has been multiplied by 2 (hence an artificial increase in the explosion energy to 2.9×10^{51} erg). The dotted line is for the doubled density in ejecta

About 7 years later the remaining core produced the second explosion with $\approx 10 M_{\odot}$ and $E_{\text{kin}} \approx 7.2 \times 10^{50}$ erg. Thus, the second ejection was faster and it had sent a shock wave into the massive cloud. The light from this radiating shock we see as SN2006gy (Fig. 2). The shock is still inside the cloud, it has not yet broken out, more than one year after the explosion of SN2006gy.

SOFT X-RAY?

There were arguments against the model of a shock moving into shells from previous explosions for SN2006gy. One can estimate the temperature of the shock using standard formulas for a rarefied medium and find that it must be very high, so SN2006gy must be a powerful source of X-ray emission. *Chandra X-ray observatory* has measured the X-ray flux on 2006 Nov 14 from the direction to SN2006gy. A best fit for X-ray luminosity in (0.5–2 keV) is 1.65×10^{39} erg/s [2], but it is a few orders of magnitude lower than expected in the naive estimates.

There is no problem with this in our model: the cloud of 25 solar masses is almost transparent to the visible light of the shock because it is almost neutral, but exactly due to this reason it is fully opaque to X-ray light. A large mass lies above the shock, see Fig. 3. The zero of the mass coordinate in Fig. 3 is the inner edge of the ejecta. The radiative shock is located at $M_r \approx 6 M_{\odot}$, or $\log r \approx 15.5$.

If we look into other data on SNe IIn we see that all of them are discovered late in X-rays! See Table 2. This is no wonder in our models: the shock is buried in the material of the first ejection.




FIGURE 3. Hydrodynamic structure at 70 days as a function of radius (left panel) and of a Lagrangean mass coordinate (right panel). Black solid line is density (left Y-axis). Violet solid line is logarithm of absolute value of luminosity $|L_{40}|$ (units 10^{40} erg/s). Dotted blue line is velocity v_8 (units 10^8 cm/s), green line is logarithm of temperature T (in K), and the red line is Rosseland optical depth τ . The scale for $\log L_{40}$, v_8 , $\log T$, and τ is on the right Y-axis.

TABLE 2. Data on X-ray observations of SNe IIn [20]

SN	galaxy	distance	discovery
		Mpc	day
1986J	NGC 891	9.6	3,300
1988Z	+03-28-022	89	2,370
1994W	NGC 4041	25	1,180
1995N	-2-38-017	24	440
1998S	NGC 3877	17	678
2005ip	NGC 2096	30	490
2005kd	PGC14370	64	450

An alternate explanation for the low X-ray flux is possible. After the work of physicists on nuclear explosions in atmosphere in the 1940s and 1950s, we know that the preheating effect becomes so large in strong supercritical shocks that the *viscous jump* in pressure and density diminishes and completely *disappears*. In *radiation dominated shocks* not only the preheating effect is important; in addition the *momentum transfer* from photons to electrons (and hence to ions, via the electric field) is very large. This also destroys the viscous jump at the shock front. Imshennik and Morozov [19] have found with an accurate accounting of the photon transfer that this happens when $P_r/P_g \simeq 8.5$. This effect implies that postshock temperature may be so low that it does not attain keV range while inside the envelope. All the heat is taken away by ‘cold’ photons. Thus, the *Chandra* results [2] tell us something about the interaction of the the first pulse ejecta with the ISM, not about the main shock penetrating the pre-ejected shell, which shines

in visible light as for SN2006gy.

CONCLUSIONS

1. Radiating shocks are the most probable sources of light in most luminous supernovae like SN2006gy.
2. The medium in which the shining shock propagates is naturally produced in massive star evolution due to violent non-linear pulsations when e^-e^+ pairs become appreciable in the pressure in their interiors.
3. The supercritical radiative shock must be well below the X-ray temperature. Even if the shock becomes hot, the overlaying matter absorbs X-rays for a long time.
4. If SN2006gy is a pulsational-pair-instability SN, a better understanding of mass-loss is needed. Otherwise, one has to find other evolution paths to SNe with double (or multiple) explosions.

ACKNOWLEDGMENTS

My work on SN2006gy was supported by NASA (at UCSC), and by the Russian Foundation for Basic Research and Science Schools (at ITEP) and by the grant IB7320-110996/1 of the Swiss National Science Foundation. My visit to Tokyo University and to this conference is supported by RESCEU. I am very grateful to S. Woosley and K. Nomoto for their warm hospitality at their Institutions and for fruitful collaboration, to M. Tanaka for sharing the observational data, and to L. Rudnick for valuable comments and improvement of my English text.

REFERENCES

1. E. Ofek, et al., *ApJ* **659**, L13 (2007).
2. N. Smith, et al., *ApJ* **666**, 1116 (2007).
3. E. Grasberg, D. Nadyozhin, *Soviet Astronomy Letters* **12**, 68 (1986).
4. N. Chugai, S. Blinnikov et al., *MNRAS* **352**, 1213 (2004).
5. E. Grasberg, V. Imshennik, D. Nadyozhin, D. K. *Astrophysics and Space Science* **10**, 28-51 (1971).
6. S. Woosley, A. Heger, and T. Weaver, *Rev.Mod.Phys.* **74**, 1015 - 1071 (2002).
7. K. Nomoto, et al., *American Institute of Physics Conference Series* **937**, 412 (2007).
8. H. Umeda, and K. Nomoto, *ArXiv e-prints* **707**, arXiv:0707.2598 (2007).
9. D. Nadyozhin, *Nauchnye Informatsii* **32**, 3 (1974).
10. S. Blinnikov, N. Dunina-Barkovskaya, and D. Nadyozhin, *ApJS* **106**, 171 (1996).
11. S. Woosley, S. Blinnikov, A. Heger, *Nature* **450**, 390 (2007).
12. N. Chugai and I. Danziger, *Astr. Lett.* **29**, 732 (2003).
13. A. Pastorello, et al., *Nature* **447**, 829 (2007).
14. T. Weaver and S. Woosley, *BAAS* **11**, 724 (1979).
15. Z. Barkat, G. Rakavy, and N. Sack, *Phys.Rev.Lett.* **18**, 379 - 381 (1967).
16. S. Woosley and T. Weaver, "The Physics of Supernovae", in *Radiation Hydrodynamics in Stars and Compact Objects*, IAU Colloq. 89 Proceedings, edited by D. Mihalas and K.-H. Winkler, Springer-Verlag, Lecture Notes in Physics 255, 1986, pp. 91 - 120.
17. A. Heger and S. Woosley, *ApJ* **567**, 532 - 543 (2002).
18. T. Weaver, G. Zimmerman, and S. Woosley, *ApJ* **225**, 1021 - 1029 (1978).

19. V. Imshennik and Yu. Morozov, *Zhurnal Prikladnoj Mekhaniki i Tekhnicheskoy Fiziki* **No. 2**, 8-21 (1964).
20. S. Immler, <http://lheawww.gsfc.nasa.gov/users/immler/> (2007).
21. K. Kawabata, M. Tanaka, et al. (2008), submitted to *ApJ*.